

Well Child Check-Up and Immunization Schedule (birth-5 years)

Each check-up includes a growth and weight evaluation, a full developmental assessment and a complete physical examination.

<u>Age at check-up</u>	<u>Immunizations/Screening Tests</u>
2 to 7 days	Post Hospital Exam and Weight Check follow ups as indicated (Hep B #1 if not given in hospital)
4 weeks*	Hep B #2, TB Screen
2 months	DTaP#1, Hib #1, IPV #1, (Pentacel), Prevnar #1, Rotavirus #1
4 months	DTaP #2, Hib #2, IPV #2, (Pentacel), Prevnar #2, Rotavirus #2
6 months	DTaP #3, Hib #3, IPV #3, (Pentacel), Prevnar #3, Rotavirus #3, TB Screen
9 months	Hep B #3, TB screen
12 months*	MMR #1, Varivax #1, Prevnar #4, Lead Screen, CBC, TB screen
15 months	DTaP #4, Hib #4, IPV #4 (Pentacel)
18 months	Hep A #1, TB screen
2 years	Hep A #2, Lead Screen, TB screen
2 ½ years (30 months)	TB screen
3 years	Vision screen, TB screen
4 years*	DTaP #5, IPV #5, MMR #2 and Varivax #2, Vision screen, TB Screen
5 years	Audio and vision screens, TB screen

***The 4 weeks, 12 months and 4 years exams should NOT be done before these ages.**

After 5 years, well child checks are recommended every year. Raleigh Pediatrics does require yearly checkups/physicals for patients who are taking medications for chronic problems such as Asthma and ADHD.

Abbreviations Used:

DTaP	Diphtheria, Tetanus, acellular Pertussis (Whooping Cough) (Ages 6 weeks-6 years)
Tdap	Tetanus, Diphtheria, acellular Pertussis (Ages 7+)
Hep B	Hepatitis B vaccine
Hep A	Hepatitis A vaccine
Hib	Haemophilus influenzae, type B (bacteria that can cause meningitis)
IPV	Inactivated Polio Vaccine
MMR	Measles, Mumps, Rubella,
Var	Varicella (Chicken Pox)
TB	Tuberculosis
Prevnar	Pneumococcal vaccine (bacteria that can cause meningitis)
Rotavirus	Oral vaccine against a common intestinal virus
CBC	Complete Blood Count – checks hemoglobin (blood test for anemia)

Adolescent vaccines

- Meningitis vaccine - given routinely between 11-16 years old (State law eff. 7/2015 requires for 7th grade school entry) , one booster after 16 years old before college entry
- Tdap (tetanus booster containing pertussis) - can be given after 6 years old but is usually done at 11-13 year well child check. (State law eff. 7/2015 requires for 7th grade school entry)
- HPV (human papilloma vaccine) - given routinely starting at 11 year well child check (If series started before age 15, then two doses are given 6 months apart; if series started at 15+, then three doses of the vaccine are given at 0, 2, and 6 month intervals)
- Varivax - second chickenpox booster if has not already had 2 or confirmed history of disease
- Hep A - if has not already had the 2 dose series.